

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International General Certificate of Secondary Education

## **MARK SCHEME for the October/November 2014 series**

### **0606 ADDITIONAL MATHEMATICS**

**0606/23**

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0606	23

1	(i)	$f(2)=0 \rightarrow 3(2)^3+8(2)^2-33(2)+p=0$ correct working to $p = 10$ AG method for quadratic factor $f(x) = (x-2)(3x^2+14x-5)$	M1 A1 M1 A1	factorise or solve quadratic factor = 0
	(ii)	$f(x) = (x-2)(3x-1)(x+5)$ $f(x)=0 \rightarrow x=2, -5, \frac{1}{3}$	M1 A1	
2	(i)	${}^{12}C_4 = 495$	B1	
	(ii)	${}^7C_2 \times {}^5C_2 = 21 \times 10$ $= 210$	M1 A1	
	(iii)	not K and B = ${}^6C_2 \times {}^4C_1 = 15 \times 4 = 60$ K and not B = ${}^6C_1 \times {}^4C_2 = 6 \times 6 = 36$ $60 + 36$ $96$  OR K and B = ${}^6C_1 \times {}^4C_1 = 6 \times 4 = 24$ not K and not B = ${}^6C_2 \times {}^4C_2 = 15 \times 6 = 90$ $210 - 90 - 24$ $96$	B1 B1 M1 A1  B1 B1 M1 A1	
3	(i)	$C$ is (1, 6) $D$ is (1, 6) + (12, 9) $= (13, 15)$	B1 M1 A1ft	correct completion      www
	(ii)	gradient of $CD = \frac{15-6}{13-1} \left( = \frac{3}{4} \right)$  gradient of $AB = \frac{10-2}{-2-4} \left( = \frac{8}{-6} = \frac{-4}{3} \right)$  $\frac{3}{4} \times \frac{-4}{3} = -1$ lines are perpendicular	B1ft  B1  B1	
	(iii)	$\text{area} = \frac{1}{2} \times AB \times CD = \frac{1}{2} \times 10 \times 15$  $= 75$ or array method	M1  A1	

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0606	23

4	(i) $2000 = 1000e^{a+b} \rightarrow a+b = \ln 2$ (ii) $3297 = 1000e^{2a-b} \rightarrow 2a+b = \ln 3.297$ oe (iii) Solve for one value $a = 0.5$ and $b = 0.193$ or $0.19$ (iv) $n = 10$ $P = 1000e^{5.193}$ $= \$180\,000.$	B1 M1 A1 M1 A1 M1 A1	substitution of 2, 3297 and rearrange
5	(i) $\overline{OX} = \mu(a+b)$ (ii) $\overline{RP} = b - 3a$ or $\overline{RX} = \lambda(b - 3a)$ oe $\overline{OX} = 3a + \lambda(b - 3a)$ (iii) $\overline{OX} = \overline{OX}$ and equate both coefficients $\mu = 3 - 3\lambda$ $\mu = \lambda$ $\mu = \lambda = 0.75$ $\frac{RX}{XP} = 3$ or $3:1$	B1 B1 B1 M1 A1 A1ft	$\frac{\lambda}{1-\lambda}$
6	(i) $m = 4$ equation of line is $\frac{\ln y - 39}{3^x - 9} = \frac{39 - 19}{9 - 4}$ $\ln y = 4(3^x) + 3$ (ii) $x = 0.5 \rightarrow \ln y = 4\sqrt{3} + 3 = 9.928$ $y = 20\,500$ (iii) Substitutes $y$ and rearrange for $3^x$ Solve $3^x = 1.150$ $x = 0.127$	B1 M1 A1ft M1 A1 M1 M1 A1	forms equation of line ft only on their gradient correct expression for $\ln y$

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0606	23

7	<p>(i) <math>x = \frac{2}{y} + 1 \rightarrow y = \frac{2}{x-1}</math>  <math>f^{-1}(x) = \frac{2}{x-1}</math></p> <p>(ii) <math>gf(x) = \left(\frac{2}{x} + 1\right)^2 + 2</math></p> <p>(iii) <math>fg(x) = \frac{2}{x^2 + 2} + 1</math></p> <p>(iv) <math>ff(x) = \frac{2}{\frac{2}{x} + 1} + 1 = \frac{2x}{x+2} + 1</math>  <math>= \frac{3x+2}{x+2}</math>  <math>\frac{3x+2}{x+2} = x \rightarrow x^2 - x - 2 = 0</math>  <math>(x-2)(x+1) = 0</math>  <math>x = 2</math> only</p>	<p>M1</p> <p>A1</p> <p>B2/1/0</p> <p>B2/1/0</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>any valid method</p> <p>-1 each error</p> <p>-1 each error</p> <p>correct starting expression</p> <p>correct algebra to given answer</p> <p>form and solve 3 term quadratic</p>
8	<p>(i) <math>v = C + K\sin 2t \quad C \neq 0</math>  <math>v = 5 + 6\sin 2t</math>  <math>a = 12\cos 2t</math></p> <p>(ii) <math>a = 0 \rightarrow \cos 2t = 0</math> and solve  <math>t = \frac{\pi}{4}</math> or 0.785 or 0.79  <math>v = 5 + 6\sin \frac{\pi}{2} = 11</math></p> <p>(iii) <math>v = 2 \rightarrow \sin 2t = -\frac{1}{2}</math> and solve  <math>t = \frac{7\pi}{12}</math> or 1.83–1.84  <math>a = 12\cos \frac{7\pi}{6} = -6\sqrt{3}</math> or -10.4</p>	<p>M1</p> <p>A1</p> <p>A1ft</p> <p>M1</p> <p>A1</p> <p>A1ft</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>set <math>a = 0</math> and solve for <math>t</math></p> <p>ft only on <math>K</math></p> <p>set <math>v = 2</math> and solve for <math>t</math></p>

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2014	0606	23

<p>9 (i)</p> $\frac{dy}{dx} = 4 - \frac{1}{(x-2)^2}$ $\frac{dy}{dx} = 0 \rightarrow (x-2)^2 = \frac{1}{4}$ $(4x^2 - 16x + 15 = 0)$ <p><math>x = 2.5</math> or <math>1.5</math>  <math>y = 12</math> or <math>4</math></p> $\frac{d^2y}{dx^2} = 2(x-2)^{-3}$ <p><math>x = 2.5 \rightarrow \frac{d^2y}{dx^2} &gt; 0 \rightarrow</math> minimum  <math>x = 1.5 \rightarrow \frac{d^2y}{dx^2} &lt; 0 \rightarrow</math> maximum</p> <p>(ii)</p> <p><math>x = 3 \rightarrow \frac{dy}{dx} = 3</math></p> <p>Use <math>m_1 m_2 = -1</math> for gradient normal from gradient tangent</p> <p>Eqn of normal : <math>\frac{y-13}{x-3} = -\frac{1}{3}</math></p> <p>Intersection of norm and curve</p> $14 - \frac{x}{3} = 4x + \frac{1}{x-2}$ $13x^2 - 68x + 87 = 0$ $x = \frac{29}{13} \text{ or } 2.23$	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b> <b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1ft</b></p> <p><b>M1</b> <b>DM1</b> <b>A1</b></p>	<p>solve 3 term quadratic from <math>\frac{dy}{dx} = 0</math></p> <p><math>x</math> values or 1 pair  <math>y</math> values or 1 pair</p> <p>use <math>\frac{d^2y}{dx^2}</math> with solution from <math>\frac{dy}{dx} = 0</math></p> <p>both identified      www</p> <p>must use numerical values</p> <p>equation and attempt to simplify  attempt to solve 3 term quadratic</p>
<p>10 (i)</p> $\text{LHS} = \frac{1 + \cos x + 1 - \cos x}{(1 - \cos x)(1 + \cos x)}$ $= \frac{2}{1 - \cos^2 x}$ $= \frac{2}{\sin^2 x} = \text{RHS}$ <p>(ii)</p> $2\text{cosec}^2 x = 8$ $\sin^2 x = \frac{1}{4}$ $\sin x = \pm \frac{1}{2}$ $x = 30^\circ, 150^\circ, 210^\circ, 330^\circ$	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>correct fraction</p> <p>correct evaluation</p> <p>use of <math>1 - \cos^2 x = \sin^2 x</math> and completion of fully correct proof</p> <p>identity used</p>